

Comment on “Higher Order Pancharatnam-Berry Phase and the Angular Momentum of Light”

In a recent letter, Milione *et al.* reported the Pancharatnam-Berry phase (PBP) on a higher-order Poincaré sphere (HOPS). They used two *spin-orbit* converters (SOCs) [1] to make a trajectory along the HOPS. However, the SOC actually does not make a path on the HOPS. We briefly discuss a way of acquiring a PBP made by SOC.

Figure 1(a) shows the configuration of the two SOC. The first and second SOC are rotated around the propagation (z) axis by θ_1 and θ_2 with respect to the x axis, respectively. We define the rotating frame as

$$(\xi_i \ \eta_i)^T = R_{-\theta_i} (x_i \ y_i)^T \quad (i = 1, 2), \quad (1)$$

where R_θ is a rotation matrix. In this frame, the reflection axis of the $4f$ -cylindrical lens pair (π converter) and the slow axis of the half-wave plate (HWP) in the i th SOC are chosen to be the ξ_i axis.

Here, we can divide the PBP into two PBPs made by π converters and by two HWPs. First, we discuss the former one. A π converter transforms the coordinate system as follow:

$$\begin{pmatrix} x' \\ y' \end{pmatrix} = R_{\theta_i} \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix} R_{-\theta_i} \begin{pmatrix} x \\ y \end{pmatrix} \quad (i = 1, 2). \quad (2)$$

We assume that the incident beam is collimated and written as $g(x, y, z) \begin{pmatrix} 1 & \sigma \end{pmatrix}^T \propto e^{il\phi} \begin{pmatrix} 1 & \sigma \end{pmatrix}^T$ at $z = 0$. The wavefunctions g at $z = 4f$ and $8f$ are respectively expressed as

$$g(x, y, 4f) = g(x', y', 0) \propto e^{2il\theta_1} g(x, y, 0), \quad (3)$$

$$g(x, y, 8f) = g(x', y', 4f) \propto e^{-2il\theta_2} g(x, y, 4f) \propto e^{2il(\theta_1 - \theta_2)} g(x, y, 0). \quad (4)$$

Thus, the PBP made by the π converters is $2l(\theta_1 - \theta_2) = -\Omega_f/2$, where Ω_f is subtended by the trajectory on the sphere for the π converter [2].

The intermediate wavefunction states in the π converters at point 2 ($z = 2f$) and point 4 ($z = 6f$), which are discussed in Ref. [3], are far from vector vortex states. Thus, the intermediate states in the π converters are not described on the higher-order Poincaré sphere but the sphere for a π converter (Fig. 1(b)). Although the states and the PBP on or along the sphere for π converters are not well analyzed yet, we think that the PBP is related to the PBP made by the Gouy phase shift [4] and by perfect reflection [5].

Needless to say, the latter PBP, which is made by two HWPs, is given by $-\Omega_l/2$, where Ω_l is the area enclosed by the trajectory on the conventional Poincaré sphere (Fig. 1(c)). By comparing Fig. 1(b) with Fig. 1(c), we

found that $\Omega_f = \Omega_l (\equiv \Omega)$. Therefore, we obtain the total PBP as

$$\gamma(C) = -\Omega_f/2 - \sigma\Omega_l/2 = -(l + \sigma)\Omega/2, \quad (5)$$

which agrees with the experimental results in the letter we comment on. We will report on the PBP made by q -plates, whose trajectories are truly along the HOPS elsewhere.

Masato Suzuki, Keisaku Yamane, Kazuhiko Oka, Yasunori Toda, and Ryuji Morita

Department of Applied Physics

Hokkaido University

Kita-13, Nishi-8, Kita-ku, Sapporo 060-8628, Japan

- [1] L. Allen, J. Courtial, and M. J. Padgett, *Phys. Rev. E* **60**, 7497 (1999).
- [2] M. J. Padgett and J. Courtial, *Opt. Lett.* **24**, 430 (1999).
- [3] V. Denisenko, V. Shvedov, A. S. Desyatnikov, D. N. Neshev, W. Krolikowski, A. Volyar, M. Soskin, and Y. S. Kivshar, *Opt. Express* **17**, 23374 (2009).
- [4] R. Simon and N. Mukunda, *Phys. Rev. Lett.* **70**, 880 (1993).
- [5] M. Berry, *Nature* **326**, 277 (1987); M. Kitano, T. Yabuzaki, and T. Ogawa, *Phys. Rev. Lett.* **58**, 523 (1987).




FIG. 1. (a) Configuration of the *spin-orbit* converters. CL1,2,3,4: cylindrical lenses; HWP1,2: half-wave plates. WFS and PoS are abbreviations for the wavefunction state and polarization state, respectively. (b) Sphere for a π converter and (c) the conventional Poincaré sphere. This figure is drawn for $\sigma = 1$.